arXiv:gr-qc/0011035v2 19 Mar 2001

Dynamics of a string coupled to gravitational waves II

— Perturbations propagate along an infinite Nambu-Goto string —

Kouji NAKAMURA*,
Department of Physics, Keio University, Hiyoshi Yokohama, 223-8521, Japan

Hideki ISHIHARA†
Tokyo Institute of Technology, Oh-Okayama Meguro-ku, Tokyo 152-0033, Japan

(February 7, 2008)

The perturbative modes propagating along an infinite string are investigated within the framework of the gauge invariant perturbation formalism on a spacetime containing a self-gravitating straight string with a finite thickness. These modes are not included in our previous analysis. We reconstruct the perturbation formalism to discuss these modes and solve the linearized Einstein equation within the first order with respect to the string oscillation amplitude. In the thin string case, we show that the oscillations of an infinite string must involve the propagation of cosmic string traveling wave.

PACS number(s): 04.30.Db,11.27.+d,98.80.Cq

In the simplest case, the dynamics of cosmic strings, which are topological defects associated with the symmetry breaking in unified theories [1], is idealized by the Nambu-Goto action. If the self-gravity of the string is ignored (test string case), the Nambu-Goto action admits oscillatory solutions. Then it is considered that strings emit gravitational emission by these oscillation and gradually lose their kinetic energy [2]. The precise dynamics of strings is important for the estimation of the energy of the gravitational waves from cosmic strings.

To clarify the precise dynamics of them, we considered the dynamics of an infinite self-gravitating Nambu-Goto string within the first order with respect to its oscillation amplitude in the previous paper [3]. It is shown that the string displacement is directly determined by the gravitational waves and there is no resonance in the gravitational wave scattering by a thin string. This shows an infinite string bends only when the gravitational wave is passing through the string and a string does not spontaneously oscillate without gravitational wave, in contrast to the fact that test strings can freely oscillate.

Though the analyses in [3] is important for the precise dynamics of strings, these are insufficient to conclude that there is no dynamical degree of freedom for free oscillations of an infinite self-gravitating Nambu-Goto string. Because the perturbative modes which propagate along the string with the light velocity is not included in the analyses in [3]. Since these modes describe the dynamics of a test Nambu-Goto string, one might expect that these modes contain the degree of freedom for free oscillations of a self-gravitating infinite string. Thus, these modes are crucial for the dynamics of an infinite string.

In this article, we concentrate on these modes, which describe the oscillations of an infinite string, within the first order with respect to its oscillation amplitude. We first solve the dynamics of an infinite thick string, after that, we consider the thin string situation as in [3]. We show the oscillations of an infinite thin Nambu-Goto string are directly given by the propagation of the gravitational waves. These are just cosmic string traveling waves discovered by Vachaspati [4] and Garfinkle [5].

As the background for the perturbation, we consider first a spacetime $(\mathcal{M}, g_{\mu\nu})$ containing a straight thick string [3,6]. The surface $\mathcal{S}$ of the thick string divides $\mathcal{M}$ into two regions: $\mathcal{M}_{ex}$ and $\mathcal{M}_{in}$. $\mathcal{M}_{in}$ is the 'thick' world sheet of the string. We also divide $\mathcal{M}$ into two submanifolds: $\mathcal{M} = \mathcal{M}_1 \times \mathcal{M}_2$ and assume the background metric on $\mathcal{M}$ in the form

$$ds^2 = \eta_{pq} dz^p dz^q + \gamma_{ab} dy^a dy^b. \tag{1}$$

The metric η_{pq} on $\mathcal{M}_2$ is the two dimensional Minkowski metric and the metric γ_{ab} on $\mathcal{M}_1$ is given by

$$\gamma_{ab} dy^a dy^b = \frac{dr^2}{1 - \hat{\alpha}^2 r^2} + r^2 d\phi^2 = d\rho^2 + \frac{\sin^2 \hat{\alpha}\rho}{\hat{\alpha}^2} d\phi^2,$$

on $\mathcal{M}_{in} \cap \mathcal{M}_1$, where $\hat{\alpha}^2 = \mathcal{R}/2 = 8\pi G\sigma_0$ and $\phi \in [0, 2\pi)$, and

$$\gamma_{ab} dy^a dy^b = \frac{dr^2}{(1-\alpha)^2} + r^2 d\phi^2 = d\rho^2 + (1-\alpha)^2 \rho^2 d\phi^2,$$

on $\mathcal{M}_{ex} \cap \mathcal{M}_1$. We shall use the indices $a, ..., d$ for tensors on $\mathcal{M}_1$ and $p, ..., s$ for those on $\mathcal{M}_2$. The Ricci scalar

---

[*] E-mail address : kouchan@phys-h.keio.ac.jp
[†] E-mail address : ishihara@th.phys.titech.ac.jp

curvature $\mathcal{R}$ on $\mathcal{M}_1$ is assumed to be a constant, and α is a deficit angle on $\mathcal{M}_{ex} \cap \mathcal{M}_1$. The metric (1) is a solution to the Einstein equation with the energy momentum tensor [1,3,6]:

$$T_{\mu\nu} = -\sigma\eta_{\mu\nu}, \quad \sigma = \begin{cases} \sigma_0 & \text{for} \quad (r \leq r_*), \\ 0 & \text{for} \quad (r > r_*), \end{cases} \tag{2}$$

where $\eta_{\mu\nu}$ is the four dimensional extension of η_{pq} and r_* is the circumference radius of $\mathcal{S} \cap \mathcal{M}_1$. Israel's conditions [7] at $\mathcal{S}$ yields $\alpha = 1 - \sqrt{1 - \hat{\alpha}^2 r_*^2}$.

To consider the metric and matter perturbations on the background (1), we expand the perturbative variables by the harmonics on $\mathcal{M}_2$. Since we concentrate only on the modes propagating along the string with the light velocity, we introduce

$$S := e^{-i\omega(t+\epsilon z)}, \quad \epsilon = \pm 1 \tag{3}$$

as the scalar harmonics. ϵ determines the direction of the wave propagation. For each ω and ϵ, we introduce the null vectors k_p and l_p defined by

$$k_p S = -i\hat{D}_p S, \quad l_p l^p = 0, \quad l_p k^p = -2\omega^2. \tag{4}$$

Using these null vectors, tensor fields on $\mathcal{M}_2$ are decomposed by the following harmonics:

$$\begin{array}{l} V^p_{(e1)} := ik^p S, \quad V^p_{(l1)} := il^p S, \quad T_{(e0)pq} := \frac{1}{2}\eta_{pq} S, \\ T_{(e2)pq} := -k_p k_q S, \quad T_{(l2)pq} := -l_p l_q S. \end{array} \tag{5}$$

S, $V^p_{(e1)}$, $T_{(e0)pq}$ and $T_{(e2)pq}$ are same as those in [3]. If k^p is neither null nor zero, there are independent of $V^p_{(o1)} := \epsilon^{pq}\hat{D}_q S$ and $T_{(o2)pq} := -\epsilon_{r(p}\hat{D}_{q)}\hat{D}^r S$ [8] in [3]. However, when k^p is null, $V^p_{(o1)}$ ($T_{(o2)pq}$) linearly depends on $V^p_{(e1)}$ ($T_{(e2)pq}$). This is the reason why our analyses in [3] fail to include the modes propagating along the string with the light velocity. Instead, (3) and (5) are the set of independent tensor harmonics for these modes and we expand the perturbation variables by these harmonics.

Let $h_{\mu\nu}$ be a perturbative metric and $t^\mu{}_\nu$ be a perturbed energy-momentum tensor. Using the harmonics (3) and (5), $h_{\mu\nu}$ and $t^\mu{}_\nu$ are expanded as follows:

$$\begin{aligned} h_{ab} &= \int f_{ab} S, \quad h_{ap} = \int \left\{ f_{a(l1)} V_{(l1)p} + f_{a(e1)} V_{(e1)p} \right\}, \\ h_{pq} &= \int \left\{ f_{(l2)} T_{(l2)pq} + f_{(e0)} T_{(e0)pq} + f_{(e2)} T_{(e2)pq} \right\}, \\ t^a{}_b &= \int s^a{}_b S, \quad t^a{}_p = \int \left\{ s^a_{(l1)} V_{(l1)p} + s^a_{(e1)} V_{(e1)p} \right\}, \\ t^p{}_q &= \int \left\{ s_{(l2)} T_{(l2)}{}^p{}_q + s_{(e0)} T_{(e0)}{}^p{}_q + s_{(e2)} T_{(e2)}{}^p{}_q \right\}, \end{aligned}$$

where $\int := \sum_{\epsilon=\pm 1} \int d\omega$. The expansion coefficients are tensors on $\mathcal{M}_1$. The perturbative energy momentum tensor $t^\mu{}_\nu$ has its support only on $\mathcal{M}_{in}$.

Here we consider the gauge-transformation of $h_{\mu\nu}$ and $t^\mu{}_\nu$ associated with $x^\mu \to x^\mu + \xi^\mu$. ξ^μ is expanded as

$$\xi_a := \int \zeta_a S, \quad \xi_p := \int \left\{ \zeta_{(l1)} V_{(l1)p} + \zeta_{(e1)} V_{(e1)p} \right\}. \tag{6}$$

Inspecting the gauge transformed variables $h_{\mu\nu} - \pounds_\xi g_{\mu\nu}$ and $t^\mu{}_\nu - \pounds_\xi T^\mu{}_\nu$, we find simple gauge-invariant combinations: For the metric perturbations,

$$\begin{array}{l} H := f_{(l2)}, \quad H_a := f_{a(l1)} - \dfrac{1}{4\omega^2} D_a f_{(e0)}, \\ H_{ab} := f_{ab} - 2D_{(a} X_{b)}. \end{array} \tag{7}$$

where D_a is the covariant derivative associated with γ_{ab} and $X^a := f^a_{(e1)} - \frac{1}{2} D^a f_{(e2)}$ is transformed to $X^a - \zeta^a$ by the gauge transformation. For the perturbations of $T^\mu{}_\nu$,

$$\begin{array}{r} \Sigma := 16\pi G (s_{(e0)} + 2X^a D_a \sigma), \\ V^a := 16\pi G (s^a_{(e1)} - \sigma X^a). \end{array} \tag{8}$$

(8) are same as those of $k^p k_p \neq 0$ modes in [3] and all coefficients in $t^\mu{}_\nu$ except for $s_{(e0)}$ and $s^a_{(e1)}$ are gauge invariant by themselves. Σ is the perturbation of the string energy density which is equal to the tangential tension. $-V^a/\mathcal{R}$ is the displacement perturbation.

In this paper, we consider the perturbative motion of an infinite Nambu-Goto string. We only consider Σ and V^a and drop the other coefficients in the above perturbative energy momentum tensor as discussed in [3].

In terms of the gauge invariant variables defined by (7) and (8), all components of the linearized Einstein equations on $\mathcal{M}_{in}$ are given by

$$\begin{array}{l} \Delta H_{ab} = \mathcal{R} H_{ab} + 2D_{(a} V_{b)} - \gamma_{ab} D_c V^c, \\ \Delta H_a - D^c D_a H_c - 2\omega^2 D_a H = 0, \\ \Delta H = 0, \\ D^c H_{ac} + 2\omega^2 H_a = V_a, \\ D^a H_a + 2\omega^2 H = 0, \\ H^c_c = 0, \end{array} \tag{9}$$

where $\Delta := D^a D_a$. On $\mathcal{M}_{ex}$, the linearized Einstein equations are given by setting $\mathcal{R} = 0 = V^a$ in (9). The perturbation of the divergence of $T_{\mu\nu}$ gives

$$2\omega^2 \sigma H^a = 0, \quad D_a V^a + \frac{1}{2}\Sigma = 0. \tag{10}$$

The first equation in (10) coincides with the perturbation of the equation of motion derived from the Nambu-Goto action [9]. The d'Alembertian of the string displacement in the first equation vanishes by virtue of $k_p k^p = 0$.

The global solutions of the perturbations on $\mathcal{M}$ should be constructed by matching the exterior and the interior solutions to (9) and (10) at $\mathcal{S}$. This is accomplished by the perturbed Israel's junction conditions $[\delta q_{\mu\nu}] := \delta q_{\mu\nu+} - \delta q_{\mu\nu-} = 0$ and $[\delta K^\mu{}_\nu] = 0$, where $\delta q_{\mu\nu\pm}$ and $\delta K^\mu{}_{\nu\pm}$ are the perturbed intrinsic metric and the extrinsic curvature of $\mathcal{S}$, respectively. The subscripts $\pm$ represent the variables facing to $\mathcal{M}_{ex}$ and $\mathcal{M}_{in}$, respectively.

Choosing the gauge freedom $\zeta_{(l)\pm}$, $\zeta_{(e)\pm}$ and $\zeta^a_\pm$ defined by (6) so that $f_{(e0)\pm} = f_{(e2)\pm} = 0$ and $X^a_+ = X^a_- = -V^a/\mathcal{R}$, the independent perturbative junction conditions for the perturbations at $\mathcal{S}$ are given by

$$\begin{aligned}
&[H] = [n^a D_a H] = 0, \\
&[n^a H_a] = [\tau^a H_a] = 0, \quad [\epsilon^{cd} D_c H_d] = 0, \\
&[\tau^a \tau^a H_{ab}] = [\tau^a n^b H_{ab}] = 0,
\end{aligned} \tag{11}$$

where ϵ^{cd} is a two-dimensional antisymmetric tensor on $\mathcal{M}_1$ and $n^a := (\partial/\partial\rho)^a$ and $\tau^a := (1/r)\,(\partial/\partial\phi)^a$.

We also impose that H, $n^a H_a$ and $\tau^a H_a$ are finite at both $r \to \infty$ and $r = 0$. Then, (9), (10) and (11) yield

$$H = 0 = H_a \tag{12}$$

on the whole spacetime $\mathcal{M}$. Further, we find that the solution to (9)-(10) are given by

$$\begin{aligned}
&H_{ab} = D_a D_b \Phi_{out}, \\
&\Phi_{out} = A_0 \ln\rho + \sum_{m=1}^{\infty} \left(A_m \rho^{-\frac{m}{1-\alpha}} + B_m \rho^{\frac{m}{1-\alpha}}\right) e^{im\phi}
\end{aligned} \tag{13}$$

on $\mathcal{M}_{ex}$, where A_m and B_m are arbitrary constants, and

$$\begin{aligned}
&H_{ab} = \left(D_a D_b - \frac{1}{2}\gamma_{ab}\Delta\right)\Phi_{in} - \epsilon_{c(a} D_{b)} D^c \Psi_{in}, \\
&V_a = \frac{1}{2} D_a (\Delta + \mathcal{R})\Phi_{in} + \frac{1}{2}\epsilon_{ac} D^c (\Delta + \mathcal{R})\Psi_{in}, \\
&\Sigma = -\Delta(\Delta + \mathcal{R})\Phi_{in}
\end{aligned} \tag{14}$$

on $\mathcal{M}_{in}$, without loss of generality [9]. V^a in (14) shows that arbitrary C^4 functions Φ_{in} and Ψ_{in} on $\mathcal{M}_{in}$ correspond to the irrotational and rotational part of the internal matter velocity field of the thick string, respectively.

(13) is just the cosmic string traveling wave discovered by Vachaspati [4] and Garfinkle [5]. This is the pp-wave exact solution to the vacuum Einstein equation. Imposing that $\tau^a \tau^b H_{ab}$ and $n^a \tau^b H_{ab}$ are finite at $r \to \infty$, we obtain $B_m = 0$ for $m \geq 1$ when $\alpha > 1/2$, while only B_1 may not vanish when $\alpha \leq 1/2$. The other coefficients in (13) depend on Φ_{in} and Ψ_{in} by the conditions (11).

In this article, we consider the oscillatory behavior of a thin string. "A thin string" should be regarded as a string whose thickness is much smaller than any other scales. The dynamics of a thin string is extracted from that of a thick string by ignoring its internal fluctuations. The thick string displacement is shown by the deformation of $\mathcal{S}$, i.e., $X^a = -V^a/\mathcal{R}$ on $\mathcal{S}$. X^a itself represents various deformations of $\mathcal{S}$. Among them, the only $m = 1$ mode in (13), the dipole deformation, represents the translation of $\mathcal{S}$ and it is relevant to the string displacement as discussed in [3]. Henceforce, we concentrate on the $m = 1$ mode.

Here, we show examples of the global solutions under the assumption that Φ_{in} and Ψ_{in} are eigen functions of Δ with a eigen value λ:

$$\Delta\Phi_{in} = -\lambda\Phi_{in}, \quad \Delta\Psi_{in} = -\lambda\Psi_{in}. \tag{15}$$

Solutions on this assumption are sufficient to demonstrate how a self-gravitating thin string behaves.

When $\lambda \neq 0$, the $m = 1$ mode solutions to (15) are

$$\Phi_{in} = C_1 P^1_\nu(x) e^{i\phi}, \quad \Psi_{in} = D_1 P^1_\nu(x) e^{i\phi}, \tag{16}$$

where $x := \sqrt{1 - \hat{\alpha}^2 r^2}$, $\nu(\nu + 1) = \lambda/\hat{\alpha}^2$. When $\lambda = 0$, we may choose $\Psi_{in} = 0$ without loss of generality and Φ_{in} of $m = 1$ is given by

$$\Phi_{in} = \hat{C}_1 \left(\frac{1 - x}{1 + x}\right)^{\frac{1}{2}} e^{i\phi}. \tag{17}$$

Here, C_1, D_1 and $\hat{C}_1$ are constants. To derive (16) and (17), the regularity at $r = 0$ is imposed.

From (11), (13), (14) and (16) or (17), we obtain

$$\begin{aligned}
A_1 &= \tfrac{1}{2}\left(\tfrac{1}{\rho_*}\right)^{-\frac{1}{1-\alpha}} (C_1 - iD_1)\,\mathcal{I}^+_1(\nu, x_*), \\
B_1 &= \tfrac{1}{2}\left(\tfrac{1}{\rho_*}\right)^{\frac{1}{1-\alpha}} (C_1 + iD_1)\,\mathcal{I}^-_1(\nu, x_*)
\end{aligned} \tag{18}$$

for the $\lambda \neq 0$ case, and

$$A_1 = 0, \quad B_1 = \hat{C}_1 \left(\frac{\alpha}{2 - \alpha}\right)^{\frac{1}{2}} \rho_*^{-\frac{1}{1-\alpha}} \tag{19}$$

for the $\lambda = 0$ case, where $x_* := \sqrt{1 - \hat{\alpha}^2 r_*^2}$ and

$$\begin{aligned}
\mathcal{I}^\pm_1(\nu, x_*) := \pm\ &\sqrt{1 - x_*^2}\, P^2_\nu(x_*) \\
&+ \tfrac{1}{2}(1 \mp x_*)(2 - \nu - \nu^2) P^1_\nu(x_*).
\end{aligned}$$

The global solution (16) and (18) includes the situations in which the string oscillates without gravitational waves outside the string. When $\mathcal{I}^\pm_1 = 0$, both A_1 and B_1 may vanish while C_1 and D_1 are nonvanishing. Since V_a does not vanish in these case, these do describe the oscillations of the string. The equation $\mathcal{I}^\pm_1 = 0$ requires the relation between ν and x_*. It can be shown that there is no solution in $0 < \nu < 1$ but the solutions exist if $\nu \geq 1$.

However, these string oscillations without gravitational waves are in the thick string case. In the global solutions given by (18) or (19), the scale of the string thickness is given by the curvature radius $\sim \mathcal{R}^{-1/2}$ on $\mathcal{M}_{in} \cap \mathcal{M}_1$ with the fixed deficit angle α, while the scale of the internal fluctuations is given by $\lambda^{-1/2}$. The condition $\lambda/\mathcal{R} \sim \nu \geq 1$ for the existence of the solution to $\mathcal{I}^\pm_1 = 0$ corresponds to the situation in which the scale of the internal fluctuations is comparable to the string thickness. The string can oscillate without the gravitational wave only when such small scale internal fluctuations are allowed.

In contrast to the thick string case, "a thin string case" is regarded as the situation $\nu \ll 1$ with the fixed α. As the thin string case, we take the leading order of ν in the global solutions. When $\nu \to 0$, $\mathcal{I}_1^{\pm}(\nu, x_*)$ behave

$$\begin{aligned} &\mathcal{I}_1^{+}(\nu, x_*) = O(\nu^2), \\ &\mathcal{I}_1^{-}(\nu, x_*) = 2\nu\left(\frac{1-x_*}{1+x_*}\right)^{\frac{1}{2}} + O(\nu^2). \end{aligned} \tag{20}$$

and all solutions (16) with (18) are reduced to a solution (17) with (19) by the replacement $\hat{C}_1 = \nu C_1$. Thus, (17) with (19) is the unique solution describes the thin string oscillations. This also shows that the traveling wave B_1 in (13) is not concerned with the internal fluctuation, while the traveling wave A_m in are excited by the internal fluctuations of the matter field.

In the same order of ν, the string displacement is

$$\begin{aligned} X^a_S &:= \int X^a|_{\mathcal{S},m=1,\lambda=0} S = -\frac{1}{2}\int SD^a\Phi_{out}|_{\mathcal{S},m=1,\lambda=0} \\ &= -\int \frac{B_1}{2(1-\alpha)}\left(\frac{r_*}{1-\alpha}\right)^{\frac{\alpha}{1-\alpha}} Se^{i\phi}\left(n^a + i\tau^a\right). \end{aligned} \tag{21}$$

Since X^a_S satisfy the equation of motion (10) derived from the Nambu-Goto action, X^a_S is regarded as the displacement of the thin Nambu-Goto string. X^a_S in (21) is completely determined by the cosmic string traveling wave B_1. The oscillations of a string are nothing but the propagation of the gravitational waves along it. This is our conclusion of the $k^p k_p = 0$ modes analyses.

The amplitude $|X^a_S| := \sqrt{\gamma_{ab}X^a_S X^b_S}$ is estimated as follows: Choosing the gauge freedom $\zeta_{(e)-}$ so that $f_{(e2)} = \Phi_{in}$, we obtain $h_{\mu\nu}dx^\mu dx^\nu = \int \Phi_{in} S\omega^2(dt + \epsilon dz)^2$. This shows the linear perturbation is not violated if $|\Phi_{in}\omega^2| < 1$. Since $|X^a_S|/r_* \sim |\Phi_{in}|\omega^2/(\omega r_*)^2$, $|X^a_S| > r_*$ if $\omega r_* \ll 1$. Therefore, the magnitude of the displacement may become larger than the string thickness within the linear perturbation framework if the wavelength of the traveling wave is sufficiently larger than the string thickness.

Though the global solutions discussed here are specific one, our conclusion in the thin string case will be hold in the wider class of the solutions (14). Since the eigen functions of Δ with different positive eigen values span the L^2 space, the internal fluctuations Φ_{in} and Ψ_{in} in L^2 space are decomposed by these eigen functions. Even so, in the thin string case, we neglect the fluctuations described by the eigen functions whose eigen values are comparable to or larger than the internal curvature. Then we will obtain (17) as the solution for the thin string case and will reach to the same conclusion as here.

We have seen that our conclusion in [3], which states that there is no dynamical degree of freedom of the free oscillations of an infinite Nambu-Goto string, is unchanged even if the $k^p k_p = 0$ modes are included into our consideration. Our formalism in [3] and here do not include the cylindrical static perturbations, which would be necessary for completion, but these have nothing to do with the dynamics of the string. Then, as the result of the systematic search of the dynamical perturbations, we conclude that the small amplitude oscillations of an infinite self-gravitating Nambu-Goto string are gravitational wave propagation. This is the essentially same conclusion as that for the thin spherical wall case in [10]. An infinite Nambu-Goto string bends only when gravitational waves are passing through the string worldsheet and continues to oscillate only when gravitational waves are propagating along itself.

The continuous string oscillations are the propagation of cosmic string traveling waves. As pointed out by Vachaspati [4], the traveling wave does not emit gravitational waves in the direction perpendicular to the string by itself. If the second or higher order in the string's oscillation amplitude give rise to the mode coupling of $k^p k_p = 0$ and $k^p k_p \neq 0$ modes, the string oscillations might emit the gravitational waves towards an observer far from the string. In the thin wall case, if there exists the gravitational wave solution propagating along the wall, the situation may be analogous.

We are pleased to thank Professor Minoru Omote and Professor Akio Hosoya for their continuous encouragement. We wish also to thank Professor Hideo Kodama for the valuable comments and discussion.

---

[1] A. Vilenkin and E.P.S Shellard, *"Cosmic Strings and Other Topological Defects"*, (Cambridge University Press, Cambridge, England, 1994) and the references therein. E.W. Kolb and M.S. Turner, *The Early Universe*, (Addison-Wesly Publishing Company, 1993) and the references therein.

[2] A. Vilenkin, and A.E. Everett, Phys. Rev. Lett. **48** 1867, (1982). T. Vachaspati, A.E. Everett and A. Vilenkin, Phys. Rev. **D30** 2046, (1984).

[3] K. Nakamura A. Ishibashi and H. Ishihara, Phys. Rev. **D62**, 101502(R), (2000).

[4] T. Vachaspati, Nucl. Phys. **B277**, 593, (1986).

[5] D. Garfinkle, Phys. Rev. **D41**, 1112, (1990). D. Garfinkle, T. Vachaspati, *ibid*, **D42**, 1960, (1990). V.P. Frolov, D. Garfinkle, *ibid*, **D42**, 3980, (1990).

[6] L. Marder, Proc. Roy. Soc. A, **244**, 524, (1958). L. Marder, *ibid*, **252**, 45, (1959). J.R. Gott, Ap.J. **288**, 422, (1985). W.A. Hiscock, Phys. Rev. **D31**, 3288, (1985). B. Linet, Gen. Rel. Grav. **17**, 1109, (1985). B. Boisseau, C. Charmousis and B. Linet, Phys. Rev. **D55**, 616, (1997).

[7] W. Israel, Nuovo Cimento **B44**, 1, (1966).

[8] ϵ_{pq} is a two-dimensional antisymmetric tensor on $\mathcal{M}_2$.

[9] K. Nakamura, in preparation.

[10] H. Kodama, H. Ishihara and Y. Fujiwara, Phys. Rev. **D50**, 7292, (1994). A. Ishibashi and H. Ishihara, *ibid*, **D56**, 3446, (1997).